



2013 Trial Examination

FORM VI

MATHEMATICS 2 UNIT

Thursday 1st August 2013

General Instructions

- Reading time — 5 minutes
- Writing time — 3 hours
- Write using black or blue pen.
- Board-approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

Total — 100 Marks

- All questions may be attempted.

Section I — 10 Marks

- Questions 1–10 are of equal value.
- Record your solutions to the multiple choice on the sheet provided.

Section II — 90 Marks

- Questions 11–16 are of equal value.
- All necessary working should be shown.
- Start each question in a new booklet.

Collection

- Write your candidate number on each booklet and on your multiple choice answer sheet.
- Hand in the booklets in a single well-ordered pile.
- Hand in a booklet for each question in Section II, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Place your multiple choice answer sheet inside the answer booklet for Question Eleven.
- Write your candidate number on this question paper and submit it with your answers.

Checklist

- SGS booklets — 6 per boy
- Multiple choice answer sheet
- Candidature — 98 boys

Examiner
BDD

SECTION I - Multiple Choice

Answers for this section should be recorded on the separate answer sheet handed out with this examination paper.

QUESTION ONE

The value of $\int_0^2 (6x^2 + 1) dx$ is:

- (A) 17
- (B) 18
- (C) 24
- (D) 66

QUESTION TWO

The line intersecting the x -axis at $x = -1$ and passing through the point $A(1, -4)$ is represented by which of the following equations?

- (A) $x + 2y - 1 = 0$
- (B) $x + 2y + 1 = 0$
- (C) $2x - y - 2 = 0$
- (D) $2x + y + 2 = 0$

QUESTION THREE

The quadratic equation $2x^2 + 12x - 9 = 0$ has roots α and β . The value of $\alpha^2\beta + \alpha\beta^2$ is:

- (A) -108
- (B) -27
- (C) 27
- (D) 108

QUESTION FOUR

What is the sum of the first ten terms of the series $96 - 48 + 24 - 12 + \dots$?

- (A) 63.9375
- (B) 191.8125
- (C) -32.736
- (D) 98.208

QUESTION FIVE

Which of the following does $\frac{d}{dx}(e^3)$ equal?


- (A) $3e^2$
- (B) e^3
- (C) 0
- (D) $\frac{1}{4}e^4$

QUESTION SIX

Which of the following statements is INCORRECT?

- (A) $\log a^n = n \log a$
- (B) $\log ab = \log a + \log b$
- (C) $\log(a - b) = \frac{\log a}{\log b}$
- (D) $\log e = 1$


QUESTION SEVEN



A particle's motion is described by the cubic graph above. Which of the following statements is NOT true of the particle at time $t = a$?

- (A) The particle's velocity is negative.
- (B) The particle has positive acceleration.
- (C) The particle is moving towards the origin.
- (D) The particle has returned to its initial position.

QUESTION EIGHT



The equation of the graph sketched above could be:


- (A) $y = 1 + \sin 2x$
- (B) $y = 1 - \sin 2x$
- (C) $y = 1 + 2 \sin 2x$
- (D) $y = 1 - 2 \sin x$

QUESTION NINE

Which of the following statements is NOT true of the function $y = x^4 + 4x^2$?

- (A) It is even.
- (B) It has a single stationary point at $x = 0$.
- (C) It has a single x -intercept at $x = 0$.
- (D) It has a single point of inflexion at $x = 0$.

QUESTION TEN



The diagram above shows the graph of a function $y = f(x)$. A pupil draws the graph of $y = 2 - |x|$ on the diagram in order to determine the number of solutions to the equation $f(x) = 2 - |x|$. His answer should be:

- (A) 0
- (B) 1
- (C) 2
- (D) 4

————— End of Section I —————

SECTION II - Written Response**Answers for this section should be recorded in the booklets provided.****Show all necessary working.****Start a new booklet for each question.****QUESTION ELEVEN (15 marks) Use a separate writing booklet.****Marks**

(a) Find the value of $\frac{e^x}{1+x^2}$ when $x = -3$. Give your answer correct to 3 decimal places. 1

(b) Differentiate:

(i) $y = \cos 2x$ 1

(ii) $y = \ln(3x + 1)$ 1

(iii) $y = e^{3x}$ 1

(c) Find the exact value of $\tan \frac{2\pi}{3}$. 1

(d) Rationalise the denominator of $\frac{1}{3 - \sqrt{5}}$. 1

(e) Find the following integrals:

(i) $\int (3x^2 + 4x) \, dx$ 1

(ii) $\int \frac{5}{x} \, dx$ 1

(iii) $\int (2x + 1)^5 \, dx$ 1

(f) Find the area of a sector subtending an angle of 6 radians at the centre of a circle of radius 3 cm. 1

(g) Find the one-hundredth term of the arithmetic sequence with first term 8 and common difference 3. 1

(h) Solve $2 \cos \theta - 1 = 0$, for $0 \leq \theta \leq 2\pi$. 2


(i) Draw a one-third page sketch of the parabola $x^2 = -8y$, carefully marking the focus and directrix. 2

QUESTION TWELVE (15 marks) Use a separate writing booklet.

Marks


(a) Find the equation of the tangent to $y = x^2 + 4x$ at $x = 1$. 2

(b)

2

The graph above shows the area bounded by the curve $y = e^{2x}$, the line $x = 3$ and the coordinate axes. Find the exact shaded area.

(c)



The points $A(2, 1)$, $B(4, 5)$ and $C(5, 2)$ have been marked in the coordinate plane above.


- (i) Show that the equation of the line passing through A and B is $2x - y - 3 = 0$. 2
- (ii) Determine the length of interval AB . 1
- (iii) Find the perpendicular distance from the point C to the line AB . 1
- (iv) Hence find the area of triangle ABC . 1

(d) Find the domain and range of $y = \sqrt{2x - 6}$. 2(e) Solve the quadratic inequation $x^2 + 2x - 3 < 0$. 2

QUESTION TWELVE (Continued)

(f)

2




The diagram above shows a regular hexagon joined to the radii of a sector. The side length of the hexagon is 5 cm. Find the exact perimeter of the resulting shape.

QUESTION THIRTEEN (15 marks) Use a separate writing booklet.

Marks

(a)



An island I , a buoy B and the mainland M lie on the vertices of a triangle, as in the diagram above. The distance from M to B is 5 km, from B to I is 10 km and from I to M is 8 km. The buoy is directly north of the mainland.

(i) Use the cosine rule to find $\angle MBI$, correct to the nearest minute.

2

(ii) What is the true bearing of the island from the buoy?

1


(b) Differentiate $y = \frac{e^{5x} + 1}{e^x}$.

2

QUESTION THIRTEEN (Continued)

(c) Consider the region bounded by the curve $y = \sec x$, the x -axis and the lines $x = 0$ 2 and $x = \frac{\pi}{3}$. Find the volume obtained by rotating this region about the x -axis.

(d)

1

The velocity-time graph of a particle is shown above. Find the distance travelled in the first 20 seconds.

(e) A particle travelling in one dimension has velocity function $v = 6 - 2t$, where v is in metres per second and t is in seconds. The particle is initially seven metres to the right of the origin. Assume that the positive direction is to the right.

- (i) Find the particle's acceleration function. 1
- (ii) Find the particle's displacement function. 1
- (iii) When is the particle at rest and what is its displacement at this time? 2
- (iv) Draw a one-third page sketch of the particle's displacement function, showing the intercepts with the axes and the vertex of this parabola. 2
- (v) What is the total distance travelled over the first eight seconds? 1

QUESTION FOURTEEN (15 marks) Use a separate writing booklet.

Marks

(a) Consider the function $f(x) = x^4 - 4x^3 + 5$.(i) Find the coordinates of the stationary points of $y = f(x)$.

3

(ii) Determine the nature of the stationary points.


2

(iii) Sketch the graph of $y = f(x)$, showing the stationary points and y -intercept.
You need not find any x -intercepts.

2

(b)


3



Find the area of the region shaded in the diagram above.

(c)

2

Find the value of x in the diagram above, giving a reason.(d) A point $P(x, y)$ moves so that its distance from $A(6, 1)$ is twice its distance from $B(-3, 4)$.(i) Show that the locus of P is a circle.

2

(ii) Find the centre and radius of the circle.

1

QUESTION FIFTEEN (15 marks) Use a separate writing booklet.**Marks**

(a) Find the exact value of $\cos \theta$ given that $\tan \theta = 7$ and $\sin \theta < 0$. 2

(b) (i) Show that $3x^2 + 4x + 5$ is positive definite. 1

(ii) Explain why the function $y = x^3 + 2x^2 + 5x + 7$ is always increasing. 1

(c) Prove that $\sin \theta \tan \theta + \cos \theta = \sec \theta$. 2

(d) Prove that $f(x) = \frac{2x}{x^2 + 1}$ is an odd function. 1

(e) (i) Differentiate xe^x . 1

(ii) Hence find $\int xe^x dx$. 1

(f) The rate of elimination $\frac{dQ}{dt}$ of a drug by the kidneys is given by the equation

$$\frac{dQ}{dt} = -kQ$$

where k is a constant and Q is the quantity of drug present in the blood. In this question, t is measured in minutes and Q in milligrams.

(i) Show that $Q = Q_0 e^{-kt}$ satisfies the equation $\frac{dQ}{dt} = -kQ$. 1

(ii) The initial quantity of drug present was measured to be 100 mg and at time $t = 20$ minutes, the quantity was 74 mg. Find the values of Q_0 and k . Give k correct to five decimal places and Q_0 to the nearest mg. 2

(iii) What is the initial rate of elimination of the drug? Give your answer correct to one decimal place. 1


(iv) How long is it until only half the original quantity of drug remains? Give your answer correct to the nearest minute. 2

The exam continues on the next page**Exam continues overleaf ...**

QUESTION SIXTEEN (15 marks) Use a separate writing booklet.

Marks

(a) In the diagram below, $\angle PQS = \angle PRS = \theta$ and $PQ = \ell$.



(i) Prove that $\triangle PQS \equiv \triangle PRS$. 1

(ii) Give a reason why $QS = RS$. 1

(iii) Show that $QR = 2\ell \cos \theta$. 1

(iv) Show that the area of $\triangle PQR$ is given by 1

$$A = \ell^2 \cos \theta \sin \theta.$$

(v) Use calculus to find the value of θ that gives the maximum area of $\triangle PQR$. 3

(b) A university student is planning to use a cash account containing \$50 000 to help fund his expenses. The account earns interest at 6% per annum, compounded monthly. At the end of each month interest is added to the account balance and then the student withdraws \$1500. Let A_n be the amount of money remaining in the account at the end of the n th month, following the student's withdrawal.

(i) Find an expression for A_1 . 1

(ii) Find expressions for A_2 and A_3 . 2

(iii) After how many months will the account have a balance of zero dollars?
Give your answer to the nearest month. 2

(c) Solve the following equation, for $0 \leq \theta \leq 2\pi$: 3

$$3 \sin^2 \theta + 3 \cos^2 \theta + 3 \tan^2 \theta + 3 \cot^2 \theta + 3 \sec^2 \theta + 3 \cosec^2 \theta = 29$$

 End of Section II

END OF EXAMINATION

B L A N K P A G E

The following list of standard integrals may be used:

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$



2013
Trial Examination
FORM VI
MATHEMATICS 2 UNIT
Thursday 1st August 2013

- Record your multiple choice answers by filling in the circle corresponding to your choice for each question.
- Fill in the circle completely.
- Each question has only one correct answer.

Question One

A B C D

Question Two

A B C D

Question Three

A B C D

Question Four

A B C D

Question Five

A B C D

Question Six

A B C D

Question Seven

A B C D

Question Eight

A B C D

Question Nine

A B C D

Question Ten

A B C D