


NEWCASTLE GRAMMAR SCHOOL

Student Number: _____



2011 TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION

Mathematics Extension 2

Examination Date: Wednesday 17th August

Examiner: Mr. M. Brain

General Instructions

- Reading Time - 5 minutes
- Working Time - 3 hours
- Write using a blue or black pen
- Write your student number on every booklet
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question
- Each question attempted is to be returned in a separate Writing Booklet clearly marked Questions 1 etc.
- If required, additional booklets may be requested

Total marks - 120

- Attempt Questions 1- 8
- All questions are of equal value

Question 1 (Start a new booklet)

Marks

a) Find **3**

(i) $\int \frac{dx}{x^2 - 16x + 60}$

(ii) $\int \frac{dx}{x^2 - 16x + 80}$

b) Evaluate **8**

(i) $\int_0^{\frac{\pi}{2}} \frac{1}{1 + \sin \theta} d\theta$

(ii) $\int_0^{\pi} e^x \cos x dx$

c) Use the substitution $u = x - 2$ to find $\int \frac{2x}{\sqrt{4x - x^2}} dx$ **4**

Question 2 (Start a new booklet)

Marks


- a) (i) Express $\frac{-1+i}{\sqrt{3}+i}$ in mod-arg form **3**
- (ii) Hence express $\cos \frac{7\pi}{12}$ in surd form
- b) Evaluate $\arg((2+i)\bar{w})$ given that $w = -1-3i$ **1**
- c) (i) On an Argand diagram shade the region where both $|z-(1+i)| \leq 1$ and $0 \leq \arg(z-(1+i)) \leq \frac{\pi}{4}$ **4**
- (ii) Find the sets of values of $|z|$ and $\arg z$ for the points in the shaded region
- d) z_1 and z_2 are two complex numbers such that $\frac{z_1+z_2}{z_1-z_2} = 2i$ **7**
- (i) On an Argand diagram show vectors representing z_1, z_2, z_1+z_2 and z_1-z_2
- (ii) Show that $|z_1| = |z_2|$
- (iii) If α is the angle between the vectors representing z_1 and z_2 show that $\tan \frac{\alpha}{2} = \frac{1}{2}$
- (iv) Show that $z_2 = \frac{1}{5}(3+4i)z_1$

Question 3 (Start a new booklet)

Marks

- a) A is a point outside a circle with centre O . P is a second point outside the circle such that $PT=PA$ where PT is a tangent to the circle at T . AO cuts the circle at D and C . PC cuts the circle at B . AB cuts the circle at E .

6



Copy the diagram into your answer booklet

- (i) Show that $\triangle PBT$ is similar to $\triangle PTC$
- (ii) Show that $\triangle APB$ is similar to $\triangle CPA$
- (iii) Hence show that DE is parallel to AP
- b) (i) On the same number plane sketch the graphs of $y = |x| - 2$ and $y = 4 + 3x - x^2$
- (ii) Hence, or otherwise, solve $\frac{|x| - 2}{4 + 3x - x^2} > 0$

3


Question 3 continued on next page

Question 3 continued

Marks

- c) The area between $y = \sin x$ and $y = \cos x$, from the y -axis to the point of intersection, A , is rotated about the line $y = 1$

4



- (i) Find the co-ordinates of point A
- (ii) Calculate the generated volume of revolution
- d) Solve the equation $8x^4 + 44x^3 + 54x^2 + 25x + 4 = 0$ given that it has a triple root


2

Question 4 (Start a new booklet)

Marks


a) Find all the roots of $P(x) = x^4 - 8x^3 + 39x^2 - 122x + 170$ given that $3 - i$ is one of the roots **4**

b) The line through the origin which is perpendicular to the tangent at $P\left(cp, \frac{c}{p}\right)$ to the rectangular hyperbola $xy = c^2$ meets the tangent at N . **7**



Show that the locus of N has the equation $(x^2 + y^2)^2 = 4c^2xy$

c) Give a possible equation for the graph below: **4**



Question 5 (Start a new booklet)

Marks

a) The equation of a curve is $x^2y^2 - x^2 + y^2 = 0$ **7**

(i) Show that the numerical value of y satisfies $|y| < 1$

(ii) Find the equations of the asymptotes

(iii) Show that $\frac{dy}{dx} = \frac{y^3}{x^3}$

(iv) Sketch the curve

b) Sketch the graph of each equation on a separate number plane: **8**

(i) (1) $y = \sqrt{x^2}$ (2) $y = (\sqrt{x})^2$

(ii) (1) $y = \ln(e^x)$ (2) $y = e^{\ln x}$

(iii) (1) $y = \sin(\sin^{-1} x)$ (2) $y = \sin^{-1}(\sin x)$

Question 6

(Start a new booklet)

Marks

- a) The inequality $\frac{a+b+c}{3} \geq \sqrt[3]{abc}$ is true for any real positive numbers a, b and c . Given that $a+b+c=1$ show:

5

(i) $\frac{1}{abc} \geq 27$

(ii) $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \geq 9$

(iii) $(1-a)(1-b)(1-c) \geq 8abc$

- b) (i) Show that the area, A , of a regular pentagon of side length P is given by

4

$$A = \frac{5}{2} P^2 \frac{\sin^2 54^\circ}{\sin 72^\circ}$$

- (ii) The area enclosed by $y = x^2$ and $y = 3$ is the base of a solid. Cross-sections of the solid, parallel to the x -axis, are regular pentagons with one side of the pentagon on the base of the solid. Calculate the volume of the solid, correct to one decimal place


Question 6 continued on next page

Question 6 continued

Marks

- c) In the diagram below the circle with the equation $(x-2)^2 + y^2 = 1$ is drawn. The region bounded by the circle is rotated about the line $x = 1$

6



- (i) Use the method of cylindrical shells to show that the volume, V , of the solid so formed is given by

$$V = 4\pi \int_1^3 (x-1)\sqrt{1-(x-2)^2} dx$$


- (ii) By using the substitution $x-2 = \sin \theta$ calculate the volume of the solid formed

Question 7 (Start a new booklet)

Marks

a) For the diagram below:

8



- (i) Show that the tangent at $P(a \sec \theta, b \tan \theta)$ on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ has equation $\frac{x \sec \theta}{a} - \frac{y \tan \theta}{b} - 1 = 0$
- (ii) Show that if the tangent at P is also a tangent to the circle with centre $(ae, 0)$ and radius $a\sqrt{e^2 + 1}$ then $\sec \theta = -e$
- (iii) Given that $\sec \theta = -e$, deduce that the points of contact, P and Q on the hyperbola, of the common tangents to the circle and the hyperbola are the extremities of a latus rectum ($x = -ae$) of the hyperbola and state the coordinates of P and Q
- (iv) Find the equations of the common tangents to the circle and the hyperbola and find the coordinates of their points of contact with the circle

b) (i) Show that $\frac{\sin(A+B) - \sin(A-B)}{2 \sin B} = \cos A$ 7

(ii) Hence show that

$$\cos x + \cos 3x + \cos 5x + \dots + \cos(2n-3)x + \cos(2n-1)x = \frac{\sin 2nx}{2 \sin x}$$


(iii) Hence evaluate $\int_0^{\frac{\pi}{2}} \frac{\sin 8x}{\sin x} dx$

Question 8

(Start a new booklet)

Marks

a) In the diagram below ABC is a triangle in which $AB = AC$ and $BC = 1$. D is the point on BC such that $\angle BAD = \theta$, $\angle CAD = 2\theta$ 6



(i) Letting $BD = x$ show that $\cos \theta = \frac{1-x}{2x}$

(ii) Hence show that $\frac{1}{3} < x < \frac{1}{2}$

b) Let α, β and γ be the non-zero roots of $x^3 + 3px + q = 0$ 5

(i) Obtain the monic equation which has the roots $\frac{\alpha\beta}{\gamma}$, $\frac{\alpha\gamma}{\beta}$ and $\frac{\beta\gamma}{\alpha}$

(ii) Show that if $\alpha\beta = \gamma$ then $(3p - q)^2 + q = 0$

c) (i) Prove that $\cos x = \sin\left(x + \frac{\pi}{2}\right)$ **4**

(ii) Given that $y = 5 \sin(x + \alpha)$ prove, by mathematical induction,
that $\frac{d^n y}{dx^n} = 5 \sin\left(x + \alpha + \frac{n\pi}{2}\right)$ for $n \geq 1$

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$